

Attention in young children who stutter

Attention levels in young children who stutter

Sarah Costelloe^a Steve Davis^b, Penny Cavenagh^a and Silviya P. Doneva^{a,*}

^aUniversity of Suffolk, Waterfront Building, England, United Kingdom

^bDivision of Psychology and Language Science, University College London, England, United Kingdom

* Corresponding Author:

Silviya P. Doneva,
University of Suffolk,
Department of Health Sciences,
Waterfront Building, 19 Neptune Quay,
Ipswich IP4 1QJ, UK
Email address: s.doneva2@uos.ac.uk

5. Acknowledgements

The financial support of the Dominic Barker Trust is gratefully acknowledged.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

Abstract

Purpose: The purpose of this study was to determine whether children who stutter (CWS) and children who do not stutter (CWNS) differ in terms of attentional ability. **Methods:** Participants were 40 age- and gender-matched CWS and CWNS (aged between 72 and 120 months). Attentional ability was assessed using the Test of Everyday Attention for Children (TEA-Ch; Manly, Robertson, Anderson & Nimmo-Smith, 1999), a clinical assessment battery comprising 13 attentional measures, assessing 3 areas of attention - selective attention, sustained attention and attentional switching. A low score on the assessment indicates attentional difficulty.

Results: There was an overall tendency for CWS to score lower than CWNS on all 13 TEA-Ch measures and all three attentional abilities. This difference reached statistical significance for the sustained attentional component.

Conclusion: The present study provides support for the hypothesis that there are some differences between CWS and CWNS in terms of attentional ability. The findings are interpreted within existing models of attention with regard to previous studies of attention in CWS.

Keywords: stuttering; attention; TEA-Ch, cognitive control

1 **1. Introduction**

2 Recent research has provided evidence to suggest that there is a link between developmental
3 stuttering and attention. Evidence from both direct experimental testing and parental reports
4 has shown that children who stutter (CWS) overall perform more poorly on attentional tasks
5 as compared to their non-stuttering counterparts (children who do not stutter; CWNS;
6 Anderson, Pellowski, Conture, & Kelly, 2003; Blood, Blood, Maloney, Weaver, & Shaffer,
7 2007; Eggers, De Nil, & Van den Bergh, 2012, 2013; Eggers, De Nil, & Bergh, 2010; Eggers
8 & Jansson-Verkasalo, 2017; Eichorn, Marton, & Pirutinsky, 2017; Felsenfeld, van
9 Beijsterveldt, & Boomsma, 2010; Karrass et al., 2006; Jo Kraft, Ambrose, & Chon, 2014;
10 Ntourou, Anderson, & Wagovich, 2018). Such findings have raised the possibility that
11 stuttering may not be an isolated disorder, but could constitute a part of a broader spectrum of
12 impairments, instead (Felsenfeld et al., 2010). Moreover, a very recently published adult study
13 also confirmed this relationship (Doneva, Davis, & Cavenagh, 2017).

14
15 The current work was instigated in conjunction with a longitudinal study on childhood
16 disfluency based in the UK (Cavenagh, Costelloe, Davis, & Howell, 2015). The initial
17 findings from this longitudinal study showed that a group of 40 CWS exhibited significant
18 deficits on the verbal, non-verbal and full IQ components of the Wechsler Pre-school and
19 Primary Scale of Intelligence (WPPSI-III; Wechsler, 2003) when compared to the 42 gender-
20 and 34 age-matched controls (CWNS). At the time of the assessment, informal reports from
21 the CWS in the study suggested that these children found it more difficult to maintain
22 concentration on the test. On the whole, children in the stuttering group had difficulties
23 staying focused and needed more adult direction to keep on task. Given the growing body of
24 literature suggesting that CWS might have attentional difficulties as compared to CWNS, the
25 researchers hypothesized that the reduced scores on the WPPSI-III, found in the study, were

1 due to less efficient attentional processing in the stuttering group rather than a difference in IQ
2 between the groups. The purpose of the present study was to test whether such attentional
3 differences between CWS and CWNS exist by incorporating a direct measure of attention (the
4 Test of Everyday Attention for Children, TEA-Ch; Manly, Robertson, Anderson & Nimmo-
5 Smith, 1999).

6 **1.1 Stuttering**

7 Stuttering or, as now referred to as childhood-onset fluency disorder, is characterized by
8 ‘disturbances in the normal fluency and time patterning of speech that are inappropriate for
9 the individual’s age and language skills, persist over time, and are characterized by frequent
10 and marked occurrences of one (or more) of the following: 1. Sound and syllable repetitions;
11 2. Sound prolongations of consonants as well as vowels; 3. Broken words (e.g., pauses within
12 a word); 4. Audible or silent blocking (filled or unfilled pauses in speech); 5. Circumlocutions
13 (word substitutions to avoid problematic words); 6. Words produced with an excess of
14 physical tension; 7. Monosyllabic whole-word repetitions (e.g., “I-I-I-I see him;)’ (American
15 Psychiatric Association, 2013, p. 45-46). Furthermore, the disorder usually starts between the
16 ages of two and four and is known to be between 2.4 to 5.33 times more prevalent in males as
17 compared to females (Andrews & Harris, 1964; Howell, Davis, & Williams, 2008). The
18 prevalence of the condition has been estimated at around 1% (Gordon, 2007), however,
19 importantly its incidence¹ is approximately 5%, with onsets occurring mainly at the preschool
20 age (Månsson, 2000). Finally, according to a systematic investigation by Blood, Ridenour,
21 Qualls and Hammer (2003), stuttering most often co-occurs with another speech disorder (in
22 their sample 33.5% of the children had an articulation disorder and 12.7% - a phonology
23 disorder); while from the comorbid non-speech disorders the most common were a learning
24 disability (15.2%), followed by a literacy disorder (8.2%) and an attention deficit disorder
25 (ADD; 5.9%). As the probability to outgrow the condition is highest in childhood, it is

1 important to examine the factors that are associated with the recovery and the persistence of
2 the disorder.

3 **1.2 The Attentional System**

4 Attention optimizes the processing resources of the individual by enhancing the information
5 of interest at a given moment and inhibiting the stimuli, identified as task-irrelevant at that
6 time (Lavie, Hirst, de Fockert, & Viding, 2004). It can operate either endogenously, driven by
7 the goals of the perceiver (e.g., I am looking for a friend in a crowd of people) or
8 exogenously, by the stimulus properties (e.g., a flashing light captures my attention).
9 Importantly, individuals have only limited perceptual and cognitive resources; when faced
10 with an abundance of information, they are forced to ‘choose’ what information to process
11 and what to leave out. However, rather than being a conscious choice, the latter takes place
12 due to the interplay between the amount and specifics of the presented stimuli (perceptual
13 load) and the amount of available resources one can allocate to their processing. Cognitive
14 control is a synonymous term to attentional control and executive function and
15 ‘encompass[es] a wide variety of cognitive processes such as dealing with novelty, planning,
16 using strategies, monitoring performance, using feedback to modify performance, vigilance,
17 and inhibiting irrelevant information’ (Cohen, Bayer, Jaudas, & Gollwitzer, 2007, p.12). It
18 lies at the heart of the three basic attentional abilities – selective attention/inhibitory control,
19 divided attention and attentional switching (Doneva et al., 2017).

20

21 Selective attention refers to one’s ability to focus on the target information in a task while
22 ignoring goal-irrelevant information. It relies on the individual’s cognitive control – the more
23 resources are available, the better the irrelevant information can be inhibited (Lavie et al.,
24 2004). In support, conditions, where the individual’s cognitive resources are hypothesized to
25 have diminished or be depleted, such as aging (Hedden & Gabrieli, 2004; Maylor & Lavie,

1 1998), anxiety (Eriksen & Hoffman, 1973) or lack of perceived power (Guinote, 2007) are all
2 associated with imperfect information selection.

3

4 Attentional switching, on the other hand, is an indicator of mental flexibility and adaptability
5 (Kreutzfeldt, Stephan, Sturm, Willmes, & Koch, 2015). It is often assessed with a card sorting
6 task where participants learn to sort cards according to a particular dimension (e.g., colour)
7 and then need to quickly adapt to the new rule for card sorting (e.g., card value) which seems
8 to change arbitrarily and could be worked out only from the experimenter's feedback on their
9 performance. To succeed in the task, one should have a good inhibitory control in order to
10 suppress the old rule (Kramer, Cepeda, & Cepeda, 2001). The latter has been well-illustrated
11 by studies with the children's card sorting task – The Dimensional Change Card Sort (DCCS)
12 task (Frye, Zelazo, & Palfai, 1995). Results on the DCCS indicate that while preschool
13 children demonstrate an understanding of the task rules, they have a difficulty switching rules
14 on the first consecutive trials after the rule change (Perner & Lang, 2002). However, older
15 children generally do not have such difficulties and are more flexible in their attentional
16 switching on the task (Zelazo, Frye, & Rapus, 1996). Furthermore, children with Attention
17 Deficit Hyperactivity Disorder (ADHD)ⁱⁱ had been found to be significantly better at
18 attentional switching when on medication, as compared to off-medication (Kramer et al.,
19 2001).

20

21 Finally, sustained attention refers to the ability to maintain focus on the task at hand without
22 allowing one's mind to shift to external distractors (Reason, 1984). This type of attentional
23 ability is often studied in go/no-go tasks where participants are required to respond to
24 frequently-appearing items, while withholding their response to a non-frequent target item
25 (Chan, 2001; Engle & Kane, 2003; Helton, Head, & Russell, 2011; Robertson, Manly,

1 Andrade, Baddeley, & Yiend, 1997). Failures to sustain attention (i.e., *lapses of attention*) are
2 quite common in everyday life (Reason, 1984). However, these are more frequent in some
3 populations, such as people with ADHD (Christakou et al., 2013), brain-damaged patients
4 (Johnson et al., 2007), aging individuals (Howard, Bessette-Symons, Zhang, & Hoyer, 2006)
5 and young children (Kannass & Oakes, 2008). The reason behind this is that sustained
6 attention relies on cognitive resources. Therefore, when cognitive resources are reduced due
7 to a resource-demanding task or the individual has a generally lower cognitive resource
8 capacity, they are easily distracted and less able to sustain attention (Engle & Kane, 2003;
9 Grandjean & Collette, 2011).

10 **1.3 Attention and stuttering**

11 Attentional ability is one area where CWS have been reported to perform more poorly as
12 compared to their fluent counterparts.

13

14 The latter has been identified for all three attentional abilities described above. For instance,
15 analysis on merged data (Alm, 2014; Eggers et al., 2010) revealed differences between CWS
16 and CWNS on several measures of attentional performance with CWS scoring higher on
17 hyperactivity and lower on attentional shifting, inhibitory control and perceptual sensitivity
18 (i.e., how quickly they notice changes in the environment). The latter has also been supported
19 by Eggers et al. (2013) who examined children's performance on a Go/No-Go task with
20 infrequent No-Go signals. This type of Go/No-Go task is believed to be a classic measure of
21 sustained attention (Robertson et al., 1997), and respectively, inhibitory control, since
22 withholding the No-Go response is particularly difficult if No-Go items are infrequent (Engle
23 & Kane, 2003). The results of this study revealed that CWS were significantly worse at
24 withholding their response to the No-Go targets, indicating that they could not make a good
25 use of their inhibitory attentional mechanism (Eggers et al., 2013)ⁱⁱⁱ. Moreover, as Go/No-Go

1 tasks can also serve as a measure of response criterion (Doneva & De Fockert, 2014; Helton
2 et al., 2011) it could be suggested that CWS were more impulsive and less capable of
3 attention switching as they did not adopt a new, more efficient response criterion to complete
4 the task (Eggers et al., 2012, 2013). Furthermore, a recent study that compared the attentional
5 switching abilities of preschool CWS and CWNS on the DCCS, found that the CWS group
6 exhibited greater RT slowing during the postswitch phase of the task and a possible tradeoff
7 between speed and accuracy for CWS after the switching of the rule has occurred (Eichorn et
8 al., 2017). Finally, the latter findings are also supported by studies, relying on parent-report
9 questionnaires – the most common of which is the BSQ (McDevitt & Carey, 1978). For
10 instance, Karrass et al. (2006) found that preschoolers who stutter were identified by their
11 parents as having significantly poorer attentional regulation, being more emotionally-reactive
12 and less able to control their emotions. Another BSQ study reported a significant difference in
13 their sample of 3.00 to 5.40 year-old CWS and controls on three out of the nine
14 temperamental traits, identified from the parental questionnaire (Anderson et al., 2003). These
15 were *distractibility* (e.g., whether the child stops an activity because something else catches
16 his/her attention); *adaptability* (e.g., how easily the child adjusts to changes in his/her routine)
17 and *rhythmicity* (e.g., whether the child spontaneously wakes up at usual time on weekends
18 and holidays). Taken together these studies suggest that stuttering is often associated with
19 poorer attention.

20

21 **1.4 Theoretical Basis**

22 There is strong evidence that despite being highly practiced, word production relies on the
23 amount of dedicated central processing resources and that the processes underlying linguistic
24 ability also underpin other, non-linguistic tasks (Cook & Meyer, 2008; Ferreira & Pashler,
25 2002; Roelofs & Piai, 2011). In fact, Ferreira and Pashler (2002) demonstrated that word

1 production is susceptible to dual-task interference, similarly to paradigms assessing
2 attentional ability. For example, when participants had to perform a picture-naming task in
3 conjunction with a tone-discriminating task, they experienced a classical psychological
4 refractory period (PRP) effect (Telford, 1931; Vince, 1949) – the shorter the time interval
5 between the two tasks, the slower were participants to discriminate the tones. Furthermore,
6 more specifically to stuttering, there are two theoretical models that aim to account for the
7 potential association between stuttering and poorer attentional performance: The Demands
8 and Capacities model (Adams, 1990; Starkweather, 1987; Starkweather & Givens-Ackerman,
9 1997) and Eggers et al.’s (2012, 2013) impeded inhibitory control account. In simple terms,
10 the Demands and Capacities model postulates that language production is atypically
11 demanding for PWS and thus the sufferer’s capacities (motor, linguistic, socio-emotional, or
12 cognitive) cannot meet the number of external demands that come with speech (e.g., time
13 pressure, speech continuity, communicating with negative listeners, etc.). On the other hand,
14 Eggers et al. (2012, 2013)’s suggest that stuttering is associated with aberrant monitoring
15 during linguistic processing and impeded inhibitory control. Moreover, as this model is based
16 on Levelt (1983), it assigns a key role to inhibitory control for the detection and processing of
17 speech errors that can arise at any of the three stages of language production:
18 conceptualization, formulation, and articulation, and that these are normally resolved by
19 processes of self-repair. However, importantly, neither the authors of the two models nor the
20 present study aim to suggest that poorer cognitive control causes stuttering or vice versa; this
21 question is beyond the scope of the current research. Our aim was to simply examine this
22 potential association to gain more insights about the nature of stuttering.

23

24

25

1 **1.5 The present study**

2 Past research on the hypothesized association between stuttering and attentional ability has
3 either relied on indirect measures of attentional ability such as parental reports (e.g., Eggers et
4 al., 2010; Felsenfeld et al., 2010) or has focused on a single attentional ability (e.g., dual
5 tasking in Maxfield et al., 2016) . The present investigation aimed to add to the other such
6 study which employed an attentional assessment battery to test participants' performance on
7 several attentional abilities at the same time (Eggers et al., 2012 who used the Attention
8 Network Test (ANT; Fan, McCandliss, Sommer, Raz, & Posner, 2002). We determined TEA-
9 Ch (Manly et al., 1999) most suitable for our purpose as it is a robust battery and one of the
10 few based on a theoretical model – (Posner & Petersen, 1990)'s Attention Network
11 Framework on which the ANT is also based. Furthermore, TEA-Ch also permits the study of
12 children's attentional performance at both the visual and the auditory domain, unlike its
13 counterpart, the ANT which has a separate visual and auditory versions (Roberts,
14 Summerfield, & Hall, 2006).

15
16 Findings from previous studies examining the attentional ability of CWS have been mixed –
17 while some found problems in sustained attention (e.g., Embrechts et al., 2000; Karrass et al.,
18 2006), others reported poorer attentional orienting (Eggers et al., 2012) or selective attention
19 (Eggers et al., 2013). It was, therefore, difficult to make a directional prediction on which
20 attentional ability will be compromised in our CWS sample, especially given that this test
21 had not been administered before to CWS, but only to adults who stutter (Doneva et al.,
22 2017). Thus, we hypothesised that CWS would perform less well on TEA-Ch compared to
23 age- and gender-matched controls, without making a specific prediction about which
24 attentional abilities will be affected.

25

1 **2. Method and Materials**

2 **2.1 Participants**

3 Forty participants were recruited for this investigation from a larger cohort that participated in
4 a longitudinal study on early childhood stuttering (Cavenagh et al., 2015). The study
5 population comprised two groups of children – children who stutter (CWS) and children who
6 do not stutter (CWNS).

7

8 The sample of CWS consisted of 20 participants altogether (16 boys and 4 girls) with an age
9 range of 72 to 106 months and a mean age of 90 months. For the purposes of determining the
10 children's level of fluency, the twenty minute speech recordings, obtained from all children at
11 the start of the longitudinal study, conducted by Cavenagh et al. (2015) were used. The
12 sample of CWS was determined, based on the following criteria: a) above 2% Syllables
13 stuttered (SS) in every 200-word speech sample; b) a score of 15 or above on the Stuttering
14 Severity Instrument – Fourth Edition (SSI-4; Riley, 2009). All of the selected CWS were
15 diagnosed with developmental stuttering by the the first author (SC) who is also a qualified
16 speech and language therapist (SLT). The time lapse between taking the speech recordings
17 and the administration of TEA-Ch was one month.

18

19 The CWNS group also consisted of 20 children (15 boys and 5 girls) with an age range of 72
20 to 120 months and a mean age of 93 months. These children were determined to be typically
21 fluent (not a child who stutters) for the following key reasons: (a) they had no present or prior
22 history of parent or teacher concern with regards to the child's speech fluency; (b) after the
23 analysis of their 20-minute recordings, obtained in the beginning of the study, the first author
24 (SC) who is also a qualified SLT did not note any concerns regarding atypical speech fluency.

25

1 Moreover, as performance on the TEA-Ch is sensitive to the participants' level of vision and
2 hearing abilities, it was also ensured that all children had normal hearing and normal or
3 corrected to normal-vision. These were assessed both by the parents/teachers and the first
4 author (SC) who met the children in person and performed the testing. There were no reports
5 of head injury from the parents of either the CWS or the CWNS. Finally, a written informed
6 consent was obtained from the parents of all children who took part of the study. The parents
7 were also fully debriefed about the purposes of the study at the end of the experimental
8 testing.

9 **2.2 Materials**

10 *Test of Attentional Ability*

11 TEA-Ch is a standardised normative clinical battery for children that allows the assessment of
12 different attentional capacities (Manly et al., 1999). It comprises of 9 subtests each making
13 different attentional demands but minimising the need for other skills such as memory,
14 language and comprehension. Furthermore, as one subtest can provide more than one
15 measures (e.g., speed, accuracy), TEA-Ch provides a total of 13 attentional measures (see the
16 description below for details). The assessment is suitable for participants aged 6-16 years (the
17 children in our sample were aged between 6 and 10 years) and provides age-normed scores.
18 The TEA-Ch covers three areas of attention: selective attention, sustained attention and
19 attentional switching. These have been explained below (Manly et al., 2001).

20

21 *Selective Attention*

22 (1) 'Sky Search' is a timed subtest where children have to find as many of the pairs of
23 identical spacecrafts as possible while ignoring the unmatched pairs, presented to
24 serve as distractors in this task. Both speed and accuracy are recorded in this task.
25 Additionally, to control for the effect of motor speed on visual attentional

1 selection, children also completed a motor control version of the test which was
2 identical to the first task, with the exception that all of the distractor items were
3 removed. A time-per-target score (time/targets found) was then calculated for both
4 conditions and then each child's motor control time per target score was subtracted
5 by their time per target score in the more attentionally demanding Sky Search
6 condition. This measure is called Sky Search Attention score.

7 (2) 'Map Mission' is a subtest in which children have to search a map to find as many
8 target symbols (i.e., pairs of knives and forks) as they can in one minute. The final
9 score represents the number of correctly identified targets (accuracy).

10 *Sustained Attention*

11 (1) 'Score!' is a subtest in which children have to silently count the number of tones
12 they hear and announce them at the end of each trial. The final score corresponds
13 to the number of trials in which the child gave a correct response (accuracy).

14 (2) 'Score Dual Task' (Score DT) requires children to do two tasks at the same time –
15 while counting the number of tones they hear, they also need to listen to an audio
16 news broadcast and notice what animal is mentioned in it. The score depends on
17 the number of correct tones and animal responses (accuracy).

18 (3) 'Sky Search Dual Task' (Sky Search DT) requires children to look for pairs of
19 identical spacecraft while counting the number of tones they hear. Here a
20 decrement measure was calculated – first each child's total time per target was
21 calculated, which was then divided by their proportion of correct scores. This
22 made the child's Sky Search DT score. Finally, to obtain the decrement, the child's
23 Sky Search score was subtracted from their Sky Search DT score.

24 (4) In 'Walk Don't Walk' children are presented with a sheet showing footprints on a
25 path made up of 14 squares. Children then have to 'walk' (with their pen on the

1 paper) when they hear a tone and ‘stop walking’ when they hear another. The
2 moves are made by dotting each square with a marker pen. The final score
3 corresponds to the number of correct responses given by the child (accuracy).

4 (5) ‘Code Transmission’ requires children to listen to a monotonous series of digits
5 while trying to identify a priorly specified target sequence (e.g., two fives
6 presented one after the other). Their task is to say the digit that came before the
7 target sequence. The final score corresponds to the number of digits, correctly
8 identified by the child (accuracy).

9

10 *Attentional Switching*

11 (1) In ‘Creature Counting’ children are presented with rows of creatures depicted in
12 their dens with arrows pointing up or down inserted between them. The task is to
13 count the number of creatures presented, while at the same time following the
14 arrows and count either upwards or downwards, depending on the direction of the
15 arrow. The number of correct responses and the time taken to complete the trials
16 were recorded (speed and accuracy).

17 (2) In ‘Opposite Worlds’ children are presented with a ‘path’ made up of the digits ‘1’
18 and ‘2’. There were two conditions in this task – ‘Same Worlds’ and ‘Opposite
19 Worlds’. In the ‘Same Worlds’ children had to pronounce the digit they see (e.g.,
20 ‘1’ when they see ‘1’) and then move on the next ‘step’ until they have reached the
21 end. In the ‘Opposite Worlds’ they had to pronounce the opposite digit of what they
22 see, instead (e.g., ‘1’ when they see ‘2’). The total time, corresponding to the
23 correct responses in each condition was recorded (speed).

24

1 **2.3 Procedure**

2 The TEA-Ch was administered by a qualified speech and language therapist (SC; the first
3 author) under the supervision of a qualified psychologist either in a quiet room at the child's
4 house or the child's school. No-one else was present during the testing.

5 **3. Results**

6 The 9 TEA-Ch subtests yielded 13 attentional measures (the number of measures for each
7 subtest is outlined in Materials; e.g., Sky Search has a speed, accuracy and an Attention score,
8 see above). Each of the raw scores was transformed into norm-referenced scaled scores that
9 adjust for age and gender, according to the TEA-Ch manual (Manly et al., 1999). The scaled
10 scores of the 13 measures were then entered into a 2x13 mixed Analysis of Variance
11 (ANOVA) with task measure as a within-participants factor (where each of the 13 TEA-Ch
12 measures was a level of this factor; see Table 1) and group (CWS vs CWNS) as a between-
13 participants factor. Following this, planned post-hoc comparisons were carried out to explore
14 the simple main effects of participants' group (CWS vs CWNS) at each of the 13 TEA-Ch
15 measures (See Section 3.1). Furthermore, a composite score for each of the three attentional
16 abilities was created from the scaled scores of the relevant attentional measures (See Section
17 3.2). Finally, the performance of the two groups was compared by performing a t test for each
18 attentional ability.

19 **3.1 Analysis on the TEA-Ch Measures**

20 The 2x13 mixed ANOVA analysis yielded a significant main effect of attentional measure (F
21 (12, 456) = 11.26, $p < .001$, partial eta sq = .229). The main effect of group was also
22 significant (F (1, 38) = 6.87, $p < .013$). Indeed, as it could be seen from the descriptive
23 statistics, CWNS overall had a higher score on all 13 measures, when compared to CWS.
24 However, no significant interaction emerged between attentional measure and group ($p >$
25 0.76). Still, for merely data exploration purposes, planned post-hoc analyses exploring the

1 simple main effects of participants' group at each of the 13 attentional measures were
2 performed. Although three of these were significant at the .05 level, none reached significance
3 at the Bonferroni adjusted alpha level for the 13 comparisons ($\alpha = .004$). These were the
4 Sky Search DT, Walk, Don't Walk, and Code Transmission subtests which all measure
5 participants' sustained attentional ability.

6 **Table 1.** Mean scaled scores for CWS and CWNS on the 13 measures of the TEA Ch.

7 -----
8 Table 1 about here
9 -----

10 **3.2 Types of Attentional Ability**

11 Composite scores for each of the three attentional abilities were created from the scaled scores
12 of the relevant measures. Selective attention was estimated by adding together the Sky Search
13 Attentional Score together with the Map mission score. The sustained attention composite
14 score was composed by Score! , Score DT, Walk Don't Walk and Code Transmission.
15 Finally, attentional switching was estimated by adding together Creature Counting, Same
16 Worlds and Different Worlds. CWS scored lower than CWNS for all three of the attention
17 categories measured by TEA-Ch (See Table 2). Importantly, the difference between the CWS
18 and CWNS group was significant for the sustained attention factor ($t(38) = 3.02, p < .005$).

19 **Table 2.** Mean scores for CWS and CWNS on type of attentional ability by TEA-Ch.

20 -----
21 Table 2 about here
22 -----
23 -----

24

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

4. Discussion

The present results revealed that when the two groups were compared on each of the three attentional abilities, the CWS had an overall poorer performance on all three abilities – selective attention, sustained attention and attentional switching, with the difference reaching statistical significance for the sustained attention factor. Furthermore, there was also a tendency for our group of CWS to consistently score lower on the TEA-Ch subtests compared to controls, especially on subtests tapping into sustained attention: Sky Search DT, Walk, Don't Walk and Code Transmission.

4.1 How the present findings fit with previous research

These results are consistent with recent published work reporting that stuttering is associated with attentional problems in both adults and children (e.g., Bosshardt, 2006; Doneva et al., 2017; Eggers et al., 2010, 2012, 2013; Eggers & Jansson-Verkasalo, 2017; Eichorn et al., 2017; Jo Kraft et al., 2014; Karrass et al., 2006; Maxfield et al., 2016; Ntourou et al., 2018). Our finding that the control group achieved an overall higher score on all 13 test measures, when compared to our stuttering group, is consistent with the notion that the disorder is associated with a general weakness in executive functioning/cognitive control. Furthermore, the latter seems to manifest differently depending on the specifics of the sample and the applied research method.

While some authors have suggested problems in inhibitory control/selective attention, others have instead reported poorer performance in attentional switching, sustained or divided attention. For example, no RT differences emerged in Eggers and Jansson-Verkasalo (2017) between their school-aged CWS and CWNS groups with regards to participants' attentional switching and inhibitory control performance as compared to baseline on an auditory

1 attentional set-shifting test (De Sonneville, 2009). However, importantly, only the control
2 group benefitted from the slowing down in responding, resulting in significantly higher
3 response accuracy for this group. Thus, arguably CWS were less able to adapt their response
4 style and experienced a higher switch cost (i.e., ‘mental “gear changing” necessary before
5 appropriate task-specific processes can proceed’; (Monsell, 2003, p. 135 in Eggers & Jansson-
6 Verkasalo, 2017). On the other hand, in their sample of preschoolers, Eichorn et al. (2017)
7 found CWS to be significantly slower than their fluent counterparts in the postswitch phase of
8 the DCCS task which again measures attentional switching. Furthermore, Piispala, Kallio,
9 Bloigu and Jansson-Verkasalo (2016) studied school-aged children of a very similar age to
10 Eggers and Jansson-Verkasalo (2017) on a visual Go/Nogo paradigm where they also
11 recorded event-related potentials (ERP). Surprisingly, neither the behavioural nor ERP data
12 revealed any differences between the groups in the Nogo condition^{iv}, showing no evidence of
13 abnormal inhibitory control in the CWS group. However, more generally, these studies taken
14 together with the present study all point to atypical attentional processing in CWS in terms of
15 stimulus evaluation, response selection and execution.

16

17 It is still challenging to determine why a conflicting pattern of findings is revealed by
18 different studies in terms of where the attentional difficulties lie in stuttering. As we already
19 mention, in our opinion, these are most likely due to the specifics of the test itself and the
20 participants’ age. Previous research has employed a number of instruments to examine the
21 attentional abilities of individuals who stutter which makes it difficult to make comparisons
22 between studies (Bosshardt, 2006; Doneva et al., 2017; Eggers et al., 2012, 2013; Embrechts
23 et al., 2000, Eichorn et al., 2017; Karrass et al., 2006; Maxfield et al., 2016). To our
24 knowledge only one study has previously explored the attentional abilities of individuals who
25 stutter using the Test of Everyday Attention (Doneva et al., 2017). This experiment was

1 conducted with adult participants and therefore, used the adult version of the test (Robertson,
2 Ward, Ridgeway, & Nimmo-Smith, 1996). The findings revealed that people who stutter
3 (PWS) demonstrated worse divided and visual selective attention. There was also a trend in
4 the results for the PWS group to perform more poorly on tasks measuring attentional
5 switching. Interestingly, the results also revealed a negative association between stuttering
6 severity and performance on two TEA subtests measuring visual selective attention (i.e., the
7 more severe one's stuttering was, the worse their performance on the task). One possibility is
8 that the discrepancy in findings between Doneva et al. (2017) and the present research can be
9 explained by the age of the participants – sustained attention is normally more problematic in
10 childhood, before consistently improving during adolescence (Klenberg, Korkman, & Lahti-
11 Nuutila, 2001; Rebok et al., 1997).

12

13 **4.2 Limitations and future direction**

14 It is important to note the limitations of the present investigation. On first place, the study
15 would have potentially benefitted from a larger number of participants in each group.
16 Unfortunately, this was not possible due to the high attrition rates we experienced and because
17 we wanted to match the two groups for number of participants, age and gender. Additionally,
18 this might be the reason why none of the comparisons between the groups on attentional
19 measure performance reached statistical significance. Arguably, Sky Search DT, Walk, Don't
20 Walk, and Code Transmission might have been significant at the Bonferroni corrected alpha
21 level if more participants were included. Still, we believe that the identified tendencies in the
22 data are a valuable contribution in the right direction for the better understanding of the
23 attentional abilities of CWS and PWS, in general. Another limitation of the present study is
24 that children were not screened for ADHD or any other relevant comorbidities; therefore, it is
25 possible that another condition, such as ADHD contributed to the overall poorer attentional

1 performance in some of the CWS. For example, by using a parent perception scale, Donaher
2 and Richels (2012) found that 21 out of the 36 school-age CWS in their sample were
3 identified by their parents as needing a further testing for their ADHD-like symptoms. Future
4 research should address this concern by controlling for ADHD and other of the most common
5 comorbidities that have been reported to co-occur with stuttering in some children (Blood et
6 al., 2003).

7
8 Finally, future research should explore the therapeutic benefits of attentional training on
9 speech fluency in CWS. For example, it might be helpful to assess the attentional skills of
10 young children prior to commencing speech therapy, so that the course can be adapted to suit
11 the needs of children with combined stuttering and attentional difficulties. For example, the
12 clinician can make allowances for the child with sustained attention difficulties by delivering
13 shorter therapy sessions, having regular breaks and rewarding effort to stay focused on tasks.
14 Communication targets could also include the areas where the child exhibits particular
15 difficulties in attentional control like listening and turn-taking. Furthermore, together with
16 Doneva et al. (2017), the present study provides support for the potential benefit of attentional
17 training programs, aimed at improving cognitive control, such as the Neurocognitive Joyful
18 Attentive Training Intervention (Nejati, Pouretamad, & Bahrami, 2013).

19 **4.3 Conclusion**

20 To conclude, when compared to fluent, age- and gender-matched controls, our CWS group
21 exhibited a significantly worse performance on the composite measure of sustained attentional
22 ability. Furthermore, the CWS achieved a lower score than the CWNS group on virtually
23 every single of the 13 attentional measures of TEA-Ch, although none of the differences could
24 reach statistical significance. The present findings lend support to the growing evidence that
25 stuttering is associated with an overall poorer attentional ability and have potential practical
26 implications for stuttering therapy.

1 **6. References**

- 2 Alm, P. A. (2014). Stuttering in relation to anxiety, temperament, and personality: Review
3 and analysis with focus on causality. *Journal of Fluency Disorders*, 40, 5–21.
4 doi:10.1016/j.jfludis.2014.01.004
- 5 American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental*
6 *Disorders* (Fifth Edition). American Psychiatric Association.
7 doi:10.1176/appi.books.9780890425596
- 8 Anderson, J. D., Pellowski, M. W., Conture, E. G., & Kelly, E. M. (2003). Temperamental
9 characteristics of young children who stutter. *Journal of Speech, Language, and*
10 *Hearing Research: JSLHR*, 46(5), 1221–1233.
- 11 Blood, G. W., Ridenour, V. J., Qualls, C. D., & Hammer, C. S. (2003). Co-occurring
12 disorders in children who stutter. *Journal of Communication Disorders*, 36(6), 427–
13 448.
- 14 Bosshardt, H. (2006). Cognitive processing load as a determinant of stuttering: Summary of a
15 research programme. *Clinical Linguistics & Phonetics*, 20(5), 371–385.
16 doi:10.1080/02699200500074321
- 17 Cavenagh, P., Costelloe, S., Davis, S., & Howell, P. (2015). Characteristics of Young
18 Children Close to the Onset of Stuttering. *Communication Disorders Quarterly*, 36(3),
19 162–171. doi:10.1177/1525740114549955
- 20 Chan, R. C. K. (2001). A further study on the sustained attention response to task (SART): the
21 effect of age, gender and education. *Brain Injury*, 15(9), 819–829.
22 doi:10.1080/02699050110034325
- 23 Christakou, A., Murphy, C. M., Chantiluke, K., Cubillo, A. I., Smith, A. B., Giampietro, V.,
24 ... Rubia, K. (2013). Disorder-specific functional abnormalities during sustained

- 1 attention in youth with Attention Deficit Hyperactivity Disorder (ADHD) and with
2 Autism. *Molecular Psychiatry*, 18(2), 236–244. doi:10.1038/mp.2011.185
- 3 Cohen, A.-L., Bayer, U. C., Jaudas, A., & Gollwitzer, P. M. (2007). Self-regulatory strategy
4 and executive control: implementation intentions modulate task switching and Simon
5 task performance. *Psychological Research*, 72(1), 12–26. doi:10.1007/s00426-006-
6 0074-2
- 7 Cook, A. E., & Meyer, A. S. (2008). Capacity demands of phoneme selection in word
8 production: New evidence from dual-task experiments. *Journal of Experimental*
9 *Psychology: Learning, Memory, and Cognition*, 34(4), 886–899. doi:10.1037/0278-
10 7393.34.4.886
- 11 De Sonneville, L. M. J. (2009). *Amsterdamse neuropsychologische taken* [Amsterdam
12 neuropsychological tasks]. Amsterdam, the Netherlands: Boom Test Publishers.
- 13 Donaher, J., & Richels, C. (2012). Traits of attention deficit/hyperactivity disorder in school-
14 age children who stutter. *Journal of Fluency Disorders*, 37(4), 242–252.
15 doi:10.1016/j.jfludis.2012.08.002
- 16 Doneva, S., Davis, S., & Cavenagh, P. (2017). Comparing the performance of people who
17 stutter and people who do not stutter on the Test of Everyday Attention. *Journal of*
18 *Clinical and Experimental Neuropsychology*, 1–15.
19 doi:10.1080/13803395.2017.1386162
- 20 Doneva, S. P., & De Fockert, J. W. (2014). More conservative go/no-go response criterion
21 under high working memory load. *Journal of Cognitive Psychology*, 26(1), 110–117.
22 doi:10.1080/20445911.2013.855780
- 23 Eggers, K., De Nil, L. F., & Van den Bergh, B. R. H. (2012). The Efficiency of Attentional
24 Networks in Children Who Stutter. *Journal of Speech, Language, and Hearing*
25 *Research*, 55(3), 946–959. doi:10.1044/1092-4388(2011/10-0208)

- 1 Eggers, K., De Nil, L. F., & Bergh, B. R. H. V. den. (2010). Temperament dimensions in
2 stuttering and typically developing children. *Journal of Fluency Disorders*, 35(4),
3 355–372. doi:10.1016/j.jfludis.2010.10.004
- 4 Eggers, K., De Nil, L. F., & Van den Bergh, B. R. H. (2013). Inhibitory control in childhood
5 stuttering. *Journal of Fluency Disorders*, 38(1), 1–13.
6 doi:10.1016/j.jfludis.2012.10.001
- 7 Eggers, K., & Jansson-Verkasalo, E. (2017). Auditory Attentional Set-Shifting and Inhibition
8 in Children Who Stutter. *Journal of Speech Language and Hearing Research*, 60(11),
9 3159. doi:10.1044/2017_JSLHR-S-16-0096
- 10 Eichorn, N., Marton, K., & Pirutinsky, S. (2017). Cognitive flexibility in preschool children
11 with and without stuttering disorders. *Journal of Fluency Disorders*.
12 doi:10.1016/j.jfludis.2017.11.001
- 13 Engle, R. W., & Kane, M. J. (2003). Executive Attention, Working Memory Capacity, and a
14 Two-Factor Theory of Cognitive Control. In *Psychology of Learning and Motivation*
15 (Vol. 44, pp. 145–199). Elsevier. doi:10.1016/S0079-7421(03)44005-X
- 16 Eriksen, C. W., & Hoffman, J. E. (1973). The extent of processing of noise elements during
17 selective encoding from visual displays. *Perception & Psychophysics*, 14(1), 155–160.
18 doi:10.3758/BF03198630
- 19 Fan, J., McCandliss, B. D., Sommer, T., Raz, A., & Posner, M. I. (2002). Testing the
20 Efficiency and Independence of Attentional Networks. *Journal of Cognitive*
21 *Neuroscience*, 14(3), 340–347. doi:10.1162/089892902317361886
- 22 Felsenfeld, S., van Beijsterveldt, C. E. M., & Boomsma, D. I. (2010). Attentional Regulation
23 in Young Twins With Probable Stuttering, High Nonfluency, and Typical Fluency.
24 *Journal of Speech Language and Hearing Research*, 53(5), 1147. doi:10.1044/1092-
25 4388(2010/09-0164)

- 1 Ferreira, V. S., & Pashler, H. (2002). Central bottleneck influences on the processing stages of
2 word production. *Journal of Experimental Psychology. Learning, Memory, and*
3 *Cognition*, 28(6), 1187–1199.
- 4 Frye, D., Zelazo, P. D., & Palfai, T. (1995). Theory of mind and rule-based reasoning.
5 *Cognitive Development*, 10(4), 483–527. doi:10.1016/0885-2014(95)90024-1
- 6 Gordon, N. (2007). Stuttering: incidence and causes. *Developmental Medicine & Child*
7 *Neurology*, 44(4), 278–282. doi:10.1111/j.1469-8749.2002.tb00806.x
- 8 Grandjean, J., & Collette, F. (2011). Influence of response prepotency strength, general
9 working memory resources, and specific working memory load on the ability to inhibit
10 predominant responses: A comparison of young and elderly participants. *Brain and*
11 *Cognition*, 77(2), 237–247. doi:10.1016/j.bandc.2011.08.004
- 12 Guinote, A. (2007). Power affects basic cognition: Increased attentional inhibition and
13 flexibility. *Journal of Experimental Social Psychology*, 43(5), 685–697.
14 doi:10.1016/j.jesp.2006.06.008
- 15 Hedden, T., & Gabrieli, J. D. E. (2004). Insights into the ageing mind: a view from cognitive
16 neuroscience. *Nature Reviews Neuroscience*, 5(2), 87–96. doi:10.1038/nrn1323
- 17 Helton, W. S., Head, J., & Russell, P. N. (2011). Reliable- and unreliable-warning cues in the
18 Sustained Attention to Response Task. *Experimental Brain Research*, 209(3), 401–
19 407. doi:10.1007/s00221-011-2563-9
- 20 Howard, M. W., Bessette-Symons, B., Zhang, Y., & Hoyer, W. J. (2006). Aging selectively
21 impairs recollection in recognition memory for pictures: Evidence from modeling and
22 receiver operating characteristic curves. *Psychology and Aging*, 21(1), 96–106.
23 doi:10.1037/0882-7974.21.1.96
- 24 Howell, P., Davis, S., & Williams, R. (2008). Late Childhood Stuttering. *Journal of Speech*
25 *Language and Hearing Research*, 51(3), 669. doi:10.1044/1092-4388(2008/048)

- 1 Jo Kraft, S., Ambrose, N., & Chon, H. (2014). Temperament and Environmental
2 Contributions to Stuttering Severity in Children: The Role of Effortful Control.
3 *Seminars in Speech and Language, 35*(02), 080–094. doi:10.1055/s-0034-1371753
- 4 Johnson, K. A., Kelly, S. P., Bellgrove, M. A., Barry, E., Cox, M., Gill, M., & Robertson, I.
5 H. (2007). Response variability in Attention Deficit Hyperactivity Disorder: Evidence
6 for neuropsychological heterogeneity. *Neuropsychologia, 45*(4), 630–638.
7 doi:10.1016/j.neuropsychologia.2006.03.034
- 8 Kannass, K. N., & Oakes, L. M. (2008). The Development of Attention and Its Relations to
9 Language in Infancy and Toddlerhood. *Journal of Cognition and Development, 9*(2),
10 222–246. doi:10.1080/15248370802022696
- 11 Karrass, J., Walden, T. A., Conture, E. G., Graham, C. G., Arnold, H. S., Hartfield, K. N., &
12 Schwenk, K. A. (2006). Relation of emotional reactivity and regulation to childhood
13 stuttering. *Journal of Communication Disorders, 39*(6), 402–423.
14 doi:10.1016/j.jcomdis.2005.12.004
- 15 Klenberg, L., Korkman, M., & Lahti-Nuutila, P. (2001). Differential Development of
16 Attention and Executive Functions in 3- to 12-Year-Old Finnish Children.
17 *Developmental Neuropsychology, 20*(1), 407–428. doi:10.1207/S15326942DN2001_6
- 18 Kramer, A. F., Cepeda, N. J., & Cepeda, M. L. (2001). Methylphenidate Effects on Task-
19 Switching Performance in Attention-Deficit/Hyperactivity Disorder. *Journal of the*
20 *American Academy of Child & Adolescent Psychiatry, 40*(11), 1277–1284.
21 doi:10.1097/00004583-200111000-00007
- 22 Kreutzfeldt, M., Stephan, D. N., Sturm, W., Willmes, K., & Koch, I. (2015). The role of
23 crossmodal competition and dimensional overlap in crossmodal attention switching.
24 *Acta Psychologica, 155*, 67–76. doi:10.1016/j.actpsy.2014.12.006

- 1 Lavie, N., Hirst, A., de Fockert, J. W., & Viding, E. (2004). Load Theory of Selective
2 Attention and Cognitive Control. *Journal of Experimental Psychology: General*,
3 133(3), 339–354. doi:10.1037/0096-3445.133.3.339
- 4 Levelt, W. (1983). Monitoring and self-repair in speech. *Cognition*, 14(1), 41–104.
5 doi:10.1016/0010-0277(83)90026-4
- 6 Manly, T., Anderson, V., Nimmo-Smith, I., Turner, A., Watson, P., & Robertson, I. H.
7 (2001). The differential assessment of children’s attention: the Test of Everyday
8 Attention for Children (TEA-Ch), normative sample and ADHD performance. *Journal*
9 *of Child Psychology and Psychiatry, and Allied Disciplines*, 42(8), 1065–1081.
- 10 Månsson, H. (2000). Childhood stuttering. *Journal of Fluency Disorders*, 25(1), 47–57.
11 doi:10.1016/S0094-730X(99)00023-6
- 12 Maxfield, N. D., Olsen, W. L., Kleinman, D., Frisch, S. A., Ferreira, V. S., & Lister, J. J.
13 (2016). Attention demands of language production in adults who stutter. *Clinical*
14 *Neurophysiology*, 127(4), 1942–1960. doi:10.1016/j.clinph.2016.01.016
- 15 Maylor, E. A., & Lavie, N. (1998). The influence of perceptual load on age differences in
16 selective attention. *Psychology and Aging*, 13(4), 563–573.
- 17 McDevitt, S. C., & Carey, W. B. (1978). The measurement of temperant in 3 - 7 year old
18 children. *Journal of Child Psychology and Psychiatry*, 19(3), 245–253.
19 doi:10.1111/j.1469-7610.1978.tb00467.x
- 20 Monsell, S. (2003). Task switching. *Trends in Cognitive Sciences*, 7(3), 134–140.
- 21 Nejati, V., Pouretemad, H. R., & Bahrami, H. (2013). Attention training in rehabilitation of
22 children with developmental stuttering. *NeuroRehabilitation*, 32(2), 297–303.
23 doi:10.3233/NRE-130847

- 1 Ntourou, K., Anderson, J. D., & Wagovich, S. A. (2018). Executive function and childhood
2 stuttering: Parent ratings and evidence from a behavioral task. *Journal of Fluency*
3 *Disorders, 56*, 18–32. doi:10.1016/j.jfludis.2017.12.001
- 4 Perner, J., & Lang, B. (2002). What causes 3-year-olds' difficulty on the dimensional change
5 card sorting task? *Infant and Child Development, 11*(2), 93–105. doi:10.1002/icd.299
- 6 Piispala, J., Kallio, M., Bloigu, R., & Jansson-Verkasalo, E. (2016). Delayed N2 response in
7 Go condition in a visual Go/Nogo ERP study in children who stutter. *Journal of*
8 *Fluency Disorders, 48*, 16–26. doi:10.1016/j.jfludis.2016.02.001
- 9 Posner, M. I., & Petersen, S. E. (1990). The Attention System of the Human Brain. *Annual*
10 *Review of Neuroscience, 13*(1), 25–42. doi:10.1146/annurev.ne.13.030190.000325
- 11 Rebok, G. W., Smith, C. B., Pascualvaca, D. M., Mirsky, A. F., Anthony, B. J., & Kellam, S.
12 G. (1997). Developmental changes in attentional performance in urban children from
13 eight to thirteen years. *Child Neuropsychology, 3*(1), 28–46.
14 doi:10.1080/09297049708401366
- 15 Roberts, K. L., Summerfield, A. Q., & Hall, D. A. (2006). Presentation modality influences
16 behavioral measures of alerting, orienting, and executive control. *Journal of the*
17 *International Neuropsychological Society: JINS, 12*(4), 485–492.
- 18 Robertson, I. H., Manly, T., Andrade, J., Baddeley, B. T., & Yiend, J. (1997). 'Oops!':
19 performance correlates of everyday attentional failures in traumatic brain injured and
20 normal subjects. *Neuropsychologia, 35*(6), 747–758.
- 21 Robertson, I. H., Ward, T., Ridgeway, V., & Nimmo-Smith, I. (1996). The structure of normal
22 human attention: The Test of Everyday Attention. *Journal of the International*
23 *Neuropsychological Society: JINS, 2*(6), 525–534.
- 24 Roelofs, A., & Piai, V. (2011). Attention demands of spoken word planning: a review.
25 *Frontiers in Psychology, 2*. doi:10.3389/fpsyg.2011.00307

- 1 Telford, C. W. (1931). The refractory phase of voluntary and associative responses. *Journal*
2 *of Experimental Psychology*, *14*(1), 1–36. doi:10.1037/h0073262
- 3 Vince, M. A. (1949). Rapid response sequences and the psychological refractory period.
4 *British Journal of Psychology. General Section*, *40*(1), 23–40. doi:10.1111/j.2044-
5 8295.1949.tb00225.x
- 6 Zelazo, P. D., Frye, D., & Rapus, T. (1996). An age-related dissociation between knowing
7 rules and using them. *Cognitive Development*, *11*(1), 37–63. doi:10.1016/S0885-
8 2014(96)90027-1
- 9

1 **Endnotes**

ⁱ How many people have ever stuttered in their life.

ⁱⁱ Symptoms of inattention in ADHD include difficulty sustaining attention on tasks, especially ones requiring mental effort, failing to give close attention to detail, inability to follow instructions, difficulty organizing tasks and activities (DSM-V; American Psychiatric Association, 2013). Moreover, symptoms of hyperactivity-impulsivity include blurting out answers, excessive talking, fidgeting and an inability to wait turn and remain seated.

ⁱⁱⁱ This was also corroborated by a more recent study by Eggers and Jansson-Verkasalo (2017) who compared the IC ability of 16 Finnish school-aged CWS to 16 Finnish fluent counterparts in another computer task and found the CWS group to be less efficient at slowing down their response latency, resulting in significantly lower response accuracy.

^{iv} The CWS differed from the CWNS group only in the ERPs in the Go condition.

Table 1. Mean scaled scores for CWS and CWNS on the 13 measures of the TEA Ch.

Attentional Ability	Measure	Group	Mean Scaled Scores (SD)	P value
	Sky Search (accuracy)	CWS	9 (2.49)	> .154
		CWNS	10.2 (2.73)	
	Sky Search (speed)	CWS	7.8 (3.04)	> .275
		CWNS	8.85 (2.96)	
Selective Attention	Sky Search (attention score)	CWS	7.9 (3.71)	> .189
		CWNS	9.5 (3.85)	
	Map Mission (accuracy)	CWS	7.85 (1.73)	> .417
		CWNS	8.35 (2.11)	
	Score! (accuracy)	CWS	10.15 (3.13)	> .170
		CWNS	11.45 (2.73)	
	Sky Search DT (decrement)	CWS	8.35 (4.71)	<.05
		CWNS	11 (3.20)	
Sustained Attention	Score DT (accuracy)	CWS	10.4 (4.38)	< .102
		CWNS	12.45 (3.28)	
	Walk Don't Walk (accuracy)	CWS	6.6 (2.62)	<.04
		CWNS	8.25 (2.20)	
	Code Transmission (accuracy)	CWS	8.25 (2.67)	<.05
		CWNS	10.10 (3.03)	
	Creature Counting (accuracy)	CWS	12.35 (2.60)	> .890
		CWNS	12.45 (1.88)	
	Creature Counting (speed)	CWS	10.35 (3.50)	> .501
		CWNS	11 (2.47)	
Attentional Switching	Same Worlds (speed)	CWS	10.25 (2.97)	> .114
		CWNS	11.85 (3.28)	
	Opposite Worlds (speed)	CWS	10.4 (3.80)	> .432
		CWNS	11.35 (3.77)	

Table 2. Mean scores for CWS and CWNS on type of attentional ability by TEA-Ch.

Attentional Ability	Group	Mean	SD	P value
Selective Attention	CWS	15.75	4.61	> .189
	CWNS	17.85	5.29	
Sustained Attention	CWS	43.75	11.25	< .005
	CWNS	53.25	8.45	
Attentional Switching	CWS	43.35	8.37	> .219
	CWNS	46.65	8.34	